

INTERSECTIONS OF QUADRICS, MOMENT-ANGLE MANIFOLDS AND CONNECTED SUMS.

Samuel Gitler.
Santiago López de Medrano.

To Sylvain Cappell, on his 65th birthday.

Introduction.

The geometric topology of the generic intersection of two homogeneous coaxial quadrics in $\mathbf{R}^n$ was studied in [LdM1] where it was shown that its intersection with the unit sphere is in most cases diffeomorphic to a triple product of spheres or to the connected sum of sphere products. The proof relied heavily on a normal form for them and involved many computations. A geometric description of the group actions on them and of their polytope quotients as well as that of the homology of those manifolds was equally valid for the intersection of any number of such quadrics, but the obstacle to extending the main result for more than two turned out to be the hopeless-looking problem of finding their normal forms, close to that of classifying all simple polytopes.

The study of those intersections continued in other directions, especially those related to the projectivizations of their complex versions (known now as *LV-M manifolds*) (see [LdM2], [LdM-V], [Me], [L-N], [B], [M-V 1], [B-M] and the recent review [Me-V 2]), giving many new examples of non-algebraic complex manifolds which fiber over toric varieties.

Following this line of research, in [B-M] a deep study of LV-M manifolds included also important advances in the study of the geometric topology of the intersection of k homogeneous quadrics in their complex versions (the manifolds now called *moment-angle manifolds*). The main questions addressed were the following:

1) Whether they can always be built up from spheres by repeatedly taking products or connected sums: they produced new examples for any k which

are so, but also showed how to construct many cases which are not. Many interesting questions arose, including a specific conjecture.

2) The transition between different topological types when the generic condition is broken at some point of a deformation (*wall-crossing*).

3) A product rule of their cohomology ring (in the spirit of the description of the homology of Z given in [LdM1]) and its applications to question 1).

Meanwhile, and independently, in [D-J] essentially the same manifolds were constructed in a more abstract way, where the main objective was to study the algebraic topology of some important quotients of them called initially *toric manifolds* and now *quasitoric manifolds*. This article originated an important development through the work of many authors, and there is a vast and deep literature along those lines for which the reader is referred to [B-P]. Yet for a long time no interchange occurred between the two lines of research involving the same objects, until a small connection appeared in the final version of [B-M] and in [D-S]. In particular, it turned out that examples relevant to question 1) above were known to these authors (see [B2]), and in [B1] Baskakov had a product rule for the cohomology ring, similar but dual to that of [B-M] mentioned in 3) above. All those examples were independent and more or less simultaneous, yet both product rules followed from an earlier computation by Buchstaber and Panov of the cohomology ring¹.

One recent expression of the line of research derived from [D-J] is the article [B-B-C-G] where a far-reaching generalization is made and a general splitting formula is derived that is, in particular, a very good geometric tool for understanding the relations among the homology groups of different spaces.

This understanding turned out to be fundamental in the process of tracing a way through the abstract situation of the intersection of $k > 2$ quadrics, thus combining efficiently both approaches to the subject as expressed in [LdM1] and [B-B-C-G] to obtain many new results. Nevertheless, the present proofs of these results do not depend logically on results of [LdM1] or [B-B-C-G] and are actually more geometric than any of them, involving practically no computations.

¹In the first version of [B-M] the product rule was derived from results in [dL], while in the published version it relied on [B-P].

The results in this article follow the three paths outlined in [B-M] mentioned above, but including now all the intersections of quadrics and not only the moment-angle manifolds:

1) In Section 1 we identify very general families of manifolds that are indeed diffeomorphic to connected sums of sphere products. The main result there is Theorem 1.3 (of which the following is a simplified version), where Z denotes an intersection of quadrics and Z^J is an infinite family obtained from Z by increasing the dimension of its coordinate spaces:

Theorem. *Assume Z is of dimension $2c$ and $(c - 1)$ -connected where $c \geq 2$. Then any Z^J of dimension at least 5 is a connected sum of sphere products.*

Particular members of such families are those moment-angle manifolds for which the result was conjectured in [B-M]. A weaker result in the odd dimensional case is given in Theorem 1.4, while Theorems 1.1 and 1.2 are relative results used in the proofs of the main theorems, but which have other applications. As a byproduct of the proofs, a simpler and neater proof of the results in [LdM1] is also obtained.

2) In section 2 we give an explicit topological description of some of the transitions, mainly that of cutting off a vertex or an edge of the associated polytope (operations $Z_{\mathbf{v}}$ and $Z_{\mathbf{e}}$). We show that, under simple assumptions, these operations preserve connected sum of sphere products and adequately modified (operations $Z_{\mathbf{v}}'$ and $Z_{\mathbf{e}}'$) they can be combined with the theorems in section 1 to give new infinite families that are diffeomorphic to connected sums of sphere products, the final result along this line being:

Theorem 2.4: *If $Z = Z(P)$ is a connected sum of sphere products and is simply connected of dimension at least 5, then any manifold obtained from Z by repeatedly applying the Z^J , $Z_{\mathbf{v}}'$ and $Z_{\mathbf{e}}'$ constructions (any number of times each and in any order), as well as its corresponding moment-angle manifold, is also a connected sum of sphere products.*

The initial manifold can be any sphere, any highly connected manifold $Z(P)$ as in Theorems 1.3 and 1.4, any simply connected manifold with $m - d = 3$ of dimension at least 5 that is not a triple sphere product ([LdM1]) or

any manifold Z^J built from a polygonal Z of the prescribed dimension and connectivity.

The first results in section 2 include cases where the initial manifold is quite general and we make a similar conjecture specific for moment-angle manifolds. A proof of that conjecture under a restrictive hypothesis is enough for us to describe the topology of another important example taken from [B-M]: the manifolds associated to the truncated cube.

3) In section 3 we use our knowledge of the above example to give a proof that their cohomology rings are not isomorphic as ungraded rings. Thus we show that the product rule for the moment-angle manifolds has to be drastically modified in the general situation. We only state the modified rule, and leave the details for another publication ([B-B-C-G-LdM]).

In section 0 we recall the necessary definitions and known results and in section 2.1 some elementary topological constructions are defined and explored. In the appendix we state and prove some results about specific differentiable manifolds, which are used in sections 1 and 2.

In the process of solving these problems several new questions and conjectures have arisen. Some extensions of our results are immediate, others should be possible by continued work along the same lines. Others seem to need a whole new approach.

Acknowledgements.

This research was initiated during a discussion with Frédéric Bosio and Laurent Meersseman around the article [A-LL] and continued through fruitful conversations with them and with Tony Bahri, Martin Bendersky, Fred Cohen, Vinicio Gómez Gutiérrez, Francisco González Acuña and Alberto Verjovsky.

Part of this research was done while the second author enjoyed the warm and stimulating hospitality of the NYU Courant Institute, for which he is most grateful to Sylvain Cappell. Invitations to Princeton University by William Browder and to the CUNY Graduate Center by Martin Bendersky and Maxim Laurentiu were also very useful and encouraging for him.

0. Preliminaries.

0.1 Basic definitions.

Let $m > d > 0$ be integers and $k = m - d - 1$,

$$\Lambda_i \in R^k, i = 1, \dots, m, \quad \Lambda = (\Lambda_1, \dots, \Lambda_m)$$

If $I \subset \{1, \dots, m\}$ we will denote by Λ_I the sequence of Λ_i with $i \in I$.

We will assume the (generic) *weak hyperbolicity* condition:

If I has k or less elements then the origin is not in the convex hull of Λ_I .

Let $Z = Z(\Lambda)$ be the submanifold² of $\mathbf{R}^m$ given by the equations

$$\sum_{i=1}^n \Lambda_i x_i^2 = 0$$

$$\sum_{i=1}^n x_i^2 = 1.$$

It is immediate that this manifold is non-empty if, and only if, the origin of R^k is in the convex hull of the Λ_i .

Z is smooth, since the weak hyperbolicity condition is equivalent to the regularity of its system of equations. It is immediate that it is stably parallelizable. A simple construction (see section 1) shows that it is always the boundary of a parallelizable manifold.

Let $Z^{\mathbf{C}} = Z^{\mathbf{C}}(\Lambda)$ be the manifold defined in $\mathbf{C}^n$ by the equations:

$$\sum_{i=1}^n \Lambda_i |z_i|^2 = 0$$

$$\sum_{i=1}^n |z_i|^2 = 1.$$

² These intersections of quadrics were first studied in [W3], [Ch], [H] and [LdM1]. The manifolds $Z^{\mathbf{C}}$ were first studied in [C-K-P], [Ch] and [LdM2] where they arose (and reappeared recently, see [Ch-LdM]) in the context of dynamical systems.

Z , $Z^{\mathbf{C}}$ are essentially the same as the manifolds constructed independently in [D-J], known as *universal abelian covers* of polytopes and *moment-angle manifolds*, respectively. Z and the constructions Z_+ , Z' , Z^J described below, can be expressed in the framework of the *generalized moment-angle complexes* defined in [B-B-C-G].

Of course, $Z^{\mathbf{C}}$ is a particular (but very important) case of a Z manifold if we write its equations in real coordinates in $\mathbf{R}^{2n}$, each $\Lambda_i \in R^k$ appearing twice.

It was shown in [LdM1] that for $k = 2$, all manifolds $Z^{\mathbf{C}}$ and almost all simply connected manifolds Z are either empty, a product of three spheres or a connected sum of manifolds, each of which is a product of two spheres. We will call this type of sums *connected sum of sphere products* (implicitly implying that each summand is the product of only two spheres).

It is easy to see that this collection of manifolds is closed under products so one gets examples for high k where Z is a product of any number of spheres and any number of connected sums of sphere products, which have therefore free homology groups and easily described cohomology rings.

In [B-M] it was shown that for $k \geq 3$ the manifold $Z^{\mathbf{C}}$ may be considerably more complicated (see also [Ba2]) and that for $k \geq 5$ its homology may have a lot of torsion. But they also conjectured that under certain conditions one obtains again the same kind of connected sums. The results of this article imply that this Bosio-Meersseman conjecture is true and we obtain the same conclusion in many more cases.

0.2 Group actions and polytopes.

The manifold Z admits a $\mathbf{Z}_2^n$ action by changing the signs of the coordinates. The quotient is a simple polytope P which can be identified with the intersection of Z and the first orthant of $\mathbf{R}^m$. It follows that Z can be reconstructed from this intersection by reflecting it on all the coordinate hyperplanes.

By a simple change of coordinates $r_i = x_i^2$, this quotient can be identified with the d -dimensional convex polytope P given by

$$\sum_{i=1}^n \Lambda_i r_i = 0.$$

$$\sum_{i=1}^n r_i = 1.$$

$$r_i \geq 0.$$

The weak hyperbolicity condition is equivalent to the fact that P is a simple polytope, meaning that each vertex is exactly in d facets of P .

The facets of P are its non-empty intersections with the coordinate hyperplanes $\{r_i = 0\}$. It follows that a non-empty Z is connected if, and only if, all these intersections are non-empty, or equivalently, if P has exactly m facets, since otherwise different components of P would lie in different sides of an hyperplane $\{x_i = 0\}$. In that case $Z = Z_* \times \mathbf{Z}_2^h$ where Z_* is a connected component of Z and h is the number of those coordinate hyperplanes not touching Z .

If we introduce a new facet by intersecting P with a half space $H = \{\sum_{i=1}^n a_i r_i \geq a\}$ whose boundary does not contain any vertex of P we obtain a new simple polytope P_H . By introducing a new coordinate $r_0 = \sum_{i=1}^n a_i r_i - a$ we can put P_H in $\mathbf{R}^{n+1}$ with $r_0 \geq 0$. By simple manipulations of the equations of P_H and changes of coordinates $r'_i = b_i r_i$, $i = 0, \dots, n$ we get a system for P_H of the same form as that of P .

This implies, by induction on the number of facets, that any simple polytope can be expressed in the above form and is therefore the quotient polytope of a connected manifold Z .³

This means that the connected Z coincide with the manifolds constructed in [D-J], called by them *universal abelian covers*.

We can therefore refer to the manifold $Z(P)$, by which we mean the diffeomorphism type of any connected manifold $Z(\Lambda)$ whose quotient polytope is P .⁴

It is known that Z is $(c - 1)$ -connected if, and only if, the intersection of any c facets of P is non-empty.⁵ Following [B-M], such a P is called *dual c -neighborly* and simply *dual neighborly* if it is $[d/2]$ -dual neighborly. The B-M conjecture says that if P is even-dimensional and dual-neighborly then $Z^{\mathbf{C}}$ is a connected sum of sphere products.

³ This *Realization Theorem* is alluded to in [W3] as a general correspondence known as the *Gale diagram* of the polytope. The *perfect* correspondence between polytopes and the manifolds Z is mentioned in [LdM], but its precise statement and proof were not included after the author learned about the Gale diagram from Wall's paper. A detailed proof using the Gale Transform is given in [B-M], Theorem 0.14. The simple proof by induction on the number of facets of P is due to Vinicio Gómez Gutiérrez, [Go].

⁴One can deduce from [B-M] that combinatorially equivalent polytopes are in fact diffeomorphic as manifolds with corners, and thus there is really only one diffeomorphism type $Z(P)$. We will not be using this rigidity result since all our constructions can be given in terms of the equations and the uniqueness follows in general from the results themselves.

⁵ The homology groups of Z follow from the computations given in [LdM1] or [B-B-C-G]. An argument for the fundamental group is given in [LdM1].

The manifold $Z^{\mathbf{C}}$ admits an action of the m -torus $T^m = (S^1)^m$ by multiplication of unit complex numbers on each of the coordinates. The quotient by this action is the same simple polytope P and $Z^{\mathbf{C}}$ can be reconstructed from $P \times T^m$ by making the obvious identifications on the facets of P . The orbits of this action over a facet F correspond to the points where some coordinate z_i vanishes and are thus the points where the corresponding factor of the torus acts trivially. This implies that these submanifolds $Z^{\mathbf{C}}(F) \subset Z^{\mathbf{C}}(P)$ have codimension 2.

If P has exactly m facets, any orbit of the action of T^m on $Z^{\mathbf{C}}$ is null-homotopic in $Z^{\mathbf{C}}$. This is because it can always be deformed to an orbit over a facet of P , where one of the factors of T^m acts trivially and so the orbit collapses into an orbit of a T^{m-1} -action. This new orbit can be deformed to a T^{m-1} orbit over another facet of P where another factor acts trivially. Since any factor of T^m acts trivially on some facet we end up with a deformation of the original orbit into a point.

This proves that in that case $Z^{\mathbf{C}}$ is simply connected, since any loop in it can be moved away from the codimension 2 manifolds lying over the facets of P and so lies in the open set $\text{interior}(P) \times T^m$ and can therefore be deformed into a single orbit which in turn deforms into a point. From the same homology computations as above, $Z^{\mathbf{C}}$ is then $2c$ -connected if, and only if, P is dual c -neighborly, and therefore always actually 2-connected.

On the contrary, if P has $m - h$ facets then $Z^{\mathbf{C}} = Z_*^{\mathbf{C}} \times \mathbf{T}^h$. (For a proof see [B-M], Lemma 0.10). In that case the orbits of T^m are not null-homotopic and $Z^{\mathbf{C}}$ is not simply connected.

The 2-connected $Z^{\mathbf{C}}$ coincide with the manifolds constructed in [D-J], called now *moment-angle manifolds*. This fact follows from the realization of a polytope as the quotient of a connected manifold Z (see footnote 3).

We can therefore refer to the manifold $Z^{\mathbf{C}}(P)$, by which we mean the diffeomorphism type of any 2-connected manifold $Z^{\mathbf{C}}(\Lambda)$ whose quotient polytope is P . It was shown in [B-P] that this diffeomorphism type and the T^m action are determined by the combinatorial type of P .

1 Opening books.

Let Λ' be obtained from Λ by adding an extra Λ_1 which we interpret as the coefficient of a new extra variable x_0 , so we get the manifold Z' :

$$\Lambda_1(x_0^2 + x_1^2) + \sum_{i>1} \Lambda_i x_i^2 = 0$$

$$x_0^2 + x_1^2 + \sum_{i>1} x_i^2 = 1.$$

Let Z_+ be the intersection of Z with the half space $x_1 \geq 0$ and Z'_+ the intersection of Z' with the half space $x_0 \geq 0$. The boundary of Z'_+ is Z . This shows that Z is always the boundary of a parallelizable manifold.

Z_+ admits an action of $\mathbf{Z}_2^{n-1}$ by changing signs on all the variables except x_1 and the quotient is again P . In other words, Z_+ can be obtained from $P \times \mathbf{Z}_2^{n-1}$ by making all the identifications on the facets of P except for the one corresponding to $x_1 = 0$ (or, equivalently, by reflecting P on all the coordinate hyperplanes except $x_1 = 0$).

Consider also the manifold Z_0 which is the intersection of Z with the subspace $x_1 = 0$. Z_0 is the boundary of Z_+ .

For example, $Z_+(\Delta^n) = D^n$ that can be obtained from $\Delta^n \times \mathbf{Z}_2^n$ by making the identifications on all the faces of Δ^n but one (or, equivalently, by reflecting the spherical simplex $S^n \cap \mathbf{R}_+^{n+1}$ on all the coordinate hyperplanes but one). Also $Z_+(\Delta^n \times I)$ can be either $D^n \times S^1$ if we do not reflect it on a facet $\Delta^{n-1} \times I$ or $S^n \times D^1$ if we do not reflect it on a facet $\Delta^n \times pt$.

The quotient polytope P' of Z' is the *book* on P obtained from $P \times I$ by collapsing $P_0 \times I$ into P_0 where the latter is the facet of P corresponding to $x_1 = 0$ (that is, P_0 is the quotient polytope of Z_0).⁶


S^1 acts on Z' (rotating the coordinates (x_0, x_1)) with fixed set Z_0 . Its quotient can be identified with Z_+ . The map

$$(x_0, x_1, x_2, \dots, x_m) \mapsto (\sqrt{x_0^2 + x_1^2}, x_2, \dots, x_m)$$


is a retraction from Z' to Z_+ which restricts to the retraction from Z to Z_+

$$(x_1, x_2, \dots, x_m) \mapsto (|x_1|, x_2, \dots, x_m)$$

⁶ This construction is implicit in [LdM1] and was much used in the process of proving its main Theorem 2. The picture of the pentagonal book appears already in [W3], p.413. The name *book* was used in [Me] and [B-M], while in the literature stemming from [D-J] it is known as the Buchstaber construction, see [B-P].



Observe further that this retraction restricted to Z'_+ is homotopic to the identity: the homotopy preserves the coordinates $x_i, i \geq 2$ and folds gradually the half space $x_0 \geq 0$ of the x_0, x_1 plane into the ray $x_0 = 0, x_1 \geq 0$ preserving the distance to the origin:



So Z is the double of Z_+ and Z' is the double of Z'_+ , and Z' is the open book with with page Z_+ and trivial holonomy.⁷

The inclusion of $Z = \partial Z'_+$ into Z'_+ is homotopic to the retraction from Z to Z_+ .

The construction of Z' from Z can be generalized as follows: Let $J = (j_1, \dots, j_m)$ be a vector of positive integers. Then we construct a new configuration Λ^J obtained from Λ by repeating j_i times the coefficient Λ_i for each i and thus a new corresponding manifold Z^J . The manifold Z^J has a natural $O(j_1) \times \dots \times O(j_m)$ action whose quotient is the same simple polytope P , as well as its own $\mathbf{Z}_2^{|J|}$ action with quotient a polytope P^J .

For example, if $J = (2, 1, \dots, 1)$ we have $Z^J = Z'$. If j is a positive integer we denote by $\underline{j}$ the vector $(j, j, \dots, j)$. Thus $Z^{\underline{1}} = Z$ and $Z^{\underline{2}}$ is $Z^{\mathbf{C}}$.

We will prove that certain families of manifolds Z are connected sums of sphere products. The rough idea of the proof in each case is as follows:

(i) We show that the lower dimensional manifolds in the family, being highly connected, are connected sums of sphere products.

(ii) Then we use the geometric relation between the manifolds Z and Z' to prove the induction step that allows us to cover all the high dimensional ones.

Part (i) can be derived from the classification results about highly connected manifolds (see, among others, [Sm], [K-M], [W1], [W2]). This type of result usually involves practically all the techniques of simply-connected surgery ([Br], [K-M]) which include the number-theoretic and algebraic analysis of the middle dimensional quadratic forms in order to represent homology classes by embedded spheres with trivial normal bundle, the construction and simplification of an adequate cobordism and the use of the h -cobordism theorem.

In our case we will not use those results but will use instead a variation of the arguments used in the induction step (ii) to prove (i) directly using the simple nature of our manifolds for which those constructions are elementary.

Part (ii) follows the spirit of those classification theorems, but is considerably simpler. We use the obvious fact that if X is a simply-connected

⁷In the notation of section 2.1, $Z = \mathcal{D}(Z_+)$, $Z' = \mathcal{D}(Z'_+)$ and $Z' = \mathcal{TOP}(Z_+)$.

connected sum of sphere products, then a basis of each homology group below the top dimension can be represented by embedded spheres with trivial normal bundle. A simple geometric lemma lets us represent various combinations of them by embedded spheres with trivial normal bundle inside $X \times I$ and disjoint from each other.

Then we use the natural cobordism Z'_+ which has the crucial but simple property that all its homology comes from its boundary Z . Combining this with the previous fact we can use Theorem A1 in the appendix to show that Z'_+ is a connected sum along the boundary of trivial bundles over spheres, so the induction step follows easily for its double Z' .

We start by the induction step (Theorem 1.1) and its variation that allows us to simplify the starting point of the induction (Theorem 1.2). These are interesting by themselves since they can be used in other situations. Then we will prove our two main results (Theorems 1.3 and 1.4).

Theorem 1.1. *Assume Z is simply connected and of dimension $d \geq 5$. If Z is a connected sum of sphere products then Z' is also a connected sum of sphere products.*

Proof:

We start with a

Lemma. *If X is a manifold of dimension d and $\alpha_1, \alpha_2 \in H_i(X)$ can be represented by disjoint embedded spheres S_1, S_2 with trivial normal bundle, then if $i \leq d - 2$, $\alpha_1 + \alpha_2$ can be represented by an embedded sphere with trivial normal bundle and disjoint from S_1 and S_2 .*

Proof of the lemma: Taking disjoint tubular neighborhoods of S_1, S_2 and joining them with a thin tube, we obtain a new manifold X' diffeomorphic to the connected sum along the boundary of two copies of $S^i \times D^{d-i}$. It is enough to prove the lemma for X' . We can take for X' a standard model constructed as follows: take S'_1, S'_2 to be round spheres in $\mathbf{R}^{i+1} \subset \mathbf{R}^d$ of radius $1/2$ centered in $(1, 0, \dots, 0)$ and $(-1, 0, \dots, 0)$ together with the straight line segment σ joining $(-1/2, 0, \dots, 0)$ and $(1/2, 0, \dots, 0)$. Now consider an ϵ neighborhood of this configuration in $\mathbf{R}^d$.

One can assume that the spheres with their standard orientations in $\mathbf{R}^{i+1}$ correspond to the classes $\alpha_1, \alpha_2 \in H_i(X)$: Because if S'_i corresponded to $-\alpha_i$,

one can reflect the product $S^i \times D^{d-i}$ on the hyperplane $x_2 = 0$ of $\mathbf{R}^d$ and twist the tube around σ accordingly.⁸

Taking the intersection of this neighborhood with $\mathbf{R}^{i+1}$ we obtain a manifold whose boundary is formed by three embedded spheres with trivial normal bundle in $\mathbf{R}^{i+1}$ and therefore also in $\mathbf{R}^d$. The outside sphere (adequately oriented) represents the sum $\alpha_1 + \alpha_2$ and is disjoint from S_1 and S_2 . *QED*

To apply the lemma to the situation of the theorem, observe that, since Z is a connected sum of sphere products, all its homology groups are free and all the generators of the homology below the top dimension are represented by embedded spheres with trivial normal bundle. Furthermore, they can all be made disjoint inside $Z \times I$ by displacing them to different levels of the coordinate $t \in I$. They all have codimension at least 2 in Z since Z is simply connected, so they have codimension at least 3 in $Z \times I$. The lemma implies that not only the generators, but all elements of the homology below the top dimension can be represented by embedded spheres in $Z \times I$ with trivial normal bundle⁹ where any finite collection of them can be represented by mutually disjoint ones.

As we showed early in this section, the inclusion of $Z = \partial Z'_+$ into Z'_+ is homotopic to a retraction. It therefore induces a split epimorphism in homology¹⁰ and in the fundamental group.

So Z'_+ is simply connected and its homology is also free and is trivial in dimensions $d - 1$ and d .

A basis for the homology of Z'_+ can then be represented by mutually disjoint embedded spheres with trivial normal bundle lying in a collar neighborhood of its boundary Z . They all have codimension at least 3, so the complement of their union in Z'_+ is still simply connected.

So all the hypotheses of Theorem A1 in the appendix are fulfilled and we conclude that Z'_+ is the connected sum along the boundary of products of

⁸ When $i = n - 1$ this last step cannot be achieved. For example, in $X = S^{d-1} \times S^1$ $\alpha + \alpha$ (for α the generator of $H_{d-1}(X)$) is not representable by an embedded sphere even if we can find two disjoint spheres each representing α .

⁹This may not be true inside Z itself: in $S^{2c} \times S^{2c}$ the sum of the generators can be represented by the diagonal sphere, but its normal bundle is the tangent bundle of S^{2c} which is not trivial. No other embedded sphere representing that sum can have trivial normal bundle since its self-intersection number is 2.

¹⁰ This was proved for $k = 2$ in [LdM1] using an explicit computation of the homology groups involved in that special case. A first proof in the general case used the stable splitting in [B-B-C-G] and the relationship between the polytopes P and P' , but its generality suggested the search for a simpler geometrical argument.

the form $S^p \times D^{d+1-p}$

Finally, Z' , being the double of Z'_+ , is the connected sum of manifolds of the form $S^p \times S^{d+1-p}$. **QED.**

Observe that all we have used about Z is that it is simply connected, all its homology groups are free and every homology class below the top dimension can be represented by an embedded sphere *inside* $Z \times I$ with trivial normal bundle. And the proof gives that Z is in fact a connected sum of sphere products. So, including the full induction on the size of J , we have actually proved more:

Theorem 1.2. *Assume that Z is simply connected and of dimension $d \geq 5$ and that all its homology groups are free and every homology class below the top dimension can be represented by an embedded sphere inside $Z \times I$ with trivial normal bundle. Then Z is a connected sum of sphere products and so is Z^J for any J .*

As a first application, assume Z is of dimension $2c$ and $(c-1)$ -connected (which means that the corresponding polytope is *dual neighborly*, see section 0.2). If $c \geq 3$ then Z clearly satisfies the hypotheses of Theorem 1.2 and every Z^J is a connected sum of sphere products. For $c = 2$ we only get that Z is h -cobordant to a sum of copies of $S^2 \times S^2$. But we can apply the construction in the proof of the theorems one more step up: one obtains that Z' is the boundary of a manifold Z''_+ which deforms down to Z'_+ . In Z'_+ all the homology classes are in dimension 2 and can be represented by embedded spheres with trivial normal bundle, so the same is true for Z''_+ . Now the dimension of Z''_+ is 6 and now Theorem A1 shows again that it is a connected sum along the boundary of copies of $S^2 \times D^4$ and so its boundary Z' is a connected sum along the boundary of copies of $S^2 \times S^3$. Applying now Theorem 1.1 inductively we obtain that all higher Z^J are connected sums of products of spheres

If $c = 1$, Z has dimension 2 and is trivially a connected sum of sphere products, and so is Z' by Theorem 2.1 since P' can be obtained from a simplex by chopping off vertices. Now it is easy to see (cf. [LdM1]) that any higher Z^J has the property that all homology classes below the top dimension can be represented by embedded spheres with trivial normal bundles and as soon as they become simply connected and of dimension at least 5 (taking $J = \underline{2}$ is more than enough) they fall into the case of Theorem 1.2. So we have:

Theorem 1.3. *Assume Z is of dimension $2c$ and $(c - 1)$ -connected. Then:*

- 1) *If $c \geq 3$, Z and any Z^J are connected sums of sphere products.*
- 2) *If $c = 2$, any Z^J of dimension at least 5 is a connected sum of sphere products.*
- 3) *If $c = 1$, Z , Z' and any simply connected Z^J of dimension at least 5 is a connected sum of sphere products.*

This includes the Bosio-Meersseman conjecture ([B-M])¹¹ for $c \geq 2$ by taking the special case $J = \underline{2}$. For $c = 1$ it includes as special cases their results about moment-angle manifolds associated to polygons and polygonal books.

The result 1) should be true for any $c \geq 1$ and without any restrictions.

Our result provides also a new, neater proof of the result in [LdM1], since the pentagonal cases (i.e., when $l = 2$ in Theorem 2 of [LdM1]) can easily be handled, starting from the case where Z is the connected sum of 5 copies of $S^2 \times S^3$ which can be proved directly (and is also covered by Theorem 1.4).

In the odd dimensional case we have for the moment only weaker results: If Z is of dimension $2c + 1$ and $(c - 1)$ -connected for $c > 1$ we need to assume that all its homology groups are free, a condition that is automatic in the even dimensional case. Nevertheless, it should be mentioned that the examples of manifolds Z with torsion using the construction in [B-M] do not have the high connectivity assumed here, so that hypothesis might still be redundant.

On the other hand we have a slightly weaker conclusion because we cannot exclude the possibility of having a connected sum of sphere bundles over spheres which are stably trivial over the embedded $(c + 1)$ -spheres but not trivial. When c is odd the total space of those bundles has torsion so this possibility is excluded by our hypothesis. When c is even the only possibility is the unit tangent bundle of S^{c+1} which is trivial only for $c = 2, 6$. In this case they cannot be discarded by their torsion. Yet it is hard to see how those non-trivial bundles could have the high $\mathbf{Z}_2^m$ symmetry of our manifolds.

In any case this problem disappears for the manifold Z' by going up to the manifold Z''_+ as above¹²:

¹¹ Thus the counterexample to this conjecture announced in [A-LL] is no such. To see directly that the argument there is not correct one can use the epimorphism part of the Hurewicz theorem to check that $\pi_6(Z)$ is not as stated.

¹² Otherwise, remark that the double of a connected sum along the boundary of stably

Theorem 1.4. *Assume Z is of dimension $2c + 1$ and $(c - 1)$ -connected where $c \geq 2$ and assume further that all the homology groups of Z are free. Then:*

(i) *Z is a connected sum of stably trivial sphere bundles over spheres and if $J \neq \underline{1}$ then Z^J is a connected sum of sphere products.*

(ii) *If c is odd or if $c = 2, 6$, then Z is a connected sum of sphere products.*

The case $J = \underline{2}$ is a weak version of the Bosio-Meersseman conjecture in the odd-dimensional case.

Theorem 1.4 is definitely not true for $c = 1$: for the cube, $Z = S^1 \times S^1 \times S^1$.


In Theorems 1.3 and 1.4 the actual dimensions of the spheres can be deduced from the computation of the homology groups of the manifolds. These can be deduced from the combinatorics of the polytope as in [LdM1], [B-P] or [B-M], or by using the stable splitting proved in [B-B-C-G]. Observe that we have also proved that the halved manifolds Z'_+ are connected sums along the boundary.

A general proof that does not use the h -cobordism theorem and that allows one to actually *see* the spheres is still badly needed.

trivial D^{c+1} -bundles over S^{c+1} is a connected sum of trivial S^{c+1} -bundles over S^{c+1} .

2 Cutting corners and making ends meet.

We study now the effect on the manifold Z of cutting off parts of the polytope P of dimension d . The general idea is the following:



Cutting off a corner of P amounts to removing a number of open balls from Z thus obtaining a manifold Z_+ with boundary. To obtain the manifold corresponding to the new polytope amounts to taking another copy of Z_+ and identifying the boundaries of the two copies (consisting each of a number of spheres), which we have simulated by joining them by tubes $D^1 \times S^{d-1}$.

We obtain the same result by taking two copies of Z and performing several times the operation of removing a disk from each copy and joining their boundary spheres by a tube. Making the operation the first time amounts to taking the connected sum of the manifolds (imagine only the lower two disks and the tube). Since the resulting manifold is connected, each new time we perform the operation amounts to attaching a handle or, equivalently, taking the connected sum with $S^1 \times S^{d-1}$. Thus the final result is the connected sum of two copies of Z and a number of copies of $S^1 \times S^{d-1}$ (one less than the number of open balls removed from Z).

To make precise this construction and some of its variants we introduce now some simple topological constructions and explore the relations among them.

2.1 Doubles, Open Books and Connected sums.

We work in the smooth category. In what follows we consider many manifolds with corners and we will use implicitly and systematically the fact that they have natural smoothings.

If Q is a manifold with boundary, we denote its boundary by ∂Q as usual.

Let $\mathcal{D}Q$ be the *double* of Q :

$$\mathcal{D}Q = \partial(Q \times D^1).$$

Let $\mathcal{TOP}(Q)$ be the open book with trivial holonomy with leaf Q and binding ∂Q , that is

$$\mathcal{TOP}(Q) = Q \times S^1 \cup \partial Q \times D^2 = \partial(Q \times D^2).$$

If M is a manifold without boundary and M_{-1} is M minus an open ball, $\mathcal{D}M_{-1}$ is diffeomorphic to $M \# (-M)$. In all cases we will consider below, M admits an orientation reversing diffeomorphism, so $\mathcal{D}M_{-1}$ will be diffeomorphic to $M \# M$.

Let $\mathcal{G}M = \partial(M_{-1} \times D^2)$.¹³

So

$$\mathcal{G}M = \mathcal{TOP}(M_{-1}).$$

As a simple example,

$$\mathcal{G}(S^n) = S^{n+1}.$$

If Q, Q' are connected d -manifolds, then we will consider their connected sum $Q \# Q'$ (connecting the *interiors* of them). If Q, Q' have also connected non-empty boundaries we can take their connected sum along the boundary $Q \amalg Q'$.

We explore now the relations between all these operations:

¹³ $\mathcal{G}$ stands for *gyration* and also for González Acuña (Fico) who defined and studied this operation in [GA].

Clearly, if M and N are closed and connected and Q, Q' are connected with connected boundary then

- a) $(M \# N)_{-1} = M_{-1} \amalg N_{-1}$.
- b) $\partial(Q \amalg Q') = \partial Q \# \partial Q'$
- c) $(Q \amalg Q') \times D^1 = (Q \times D^1) \amalg (Q' \times D^1)$
- d) $\mathcal{D}(Q \amalg Q') = \mathcal{D}Q \# \mathcal{D}Q'$.

More surprising are the following facts:

Lemma 1 (Fico's Lemmata, [GA]):¹⁴

- 1) If M and N are connected, then $\mathcal{G}(M \# N) = \mathcal{G}M \# \mathcal{G}N$.
- 2) $\mathcal{G}(S^p \times S^q) = (S^{p+1} \times S^q) \# (S^p \times S^{q+1})$.

Proof: For 1), just observe from a) and c) above that

$$(M \# N)_{-1} \times D^2 = (M_{-1} \amalg N_{-1}) \times D^2 = (M_{-1} \times D^2) \amalg (N_{-1} \times D^2)$$

and take the boundary using b).

As for 2), observe that

$$(S^p \times S^q)_{-1} = (S^p \times D^q) \cup_{D^p \times D^q} (D^p \times S^q)$$

(the *plumbing* of two trivial disk bundles). This is not the connected sum $(S^p \times D^q) \amalg (D^p \times S^q)$, since its boundary is a sphere, the underlying fact being that the two core spheres cannot be separated. However, it is easy to see, by separating the spheres in the product with D^1 , that

$$((S^p \times D^q) \cup_{D^p \times D^q} (D^p \times S^q)) \times D^1 = (S^p \times D^{q+1}) \amalg (D^{p+1} \times S^q)$$

Taking again the product with D^1 , property c) and the boundary, we obtain 2).

¹⁴Unfortunately the Leibniz-Fico rule: $\mathcal{G}(M \times N) = (\mathcal{G}M \times N) \# (M \times \mathcal{G}N)$ generalizing 2) is only valid when both M and N are spheres.

Extending these ideas we have

Lemma 2. *Let Q, Q' be connected d -manifolds. Then*
(A) *If Q, Q' have non-empty boundary then*

$$\mathcal{D}(Q \# Q') = \mathcal{D}Q \# \mathcal{D}Q' \# (S^1 \times S^{d-1})$$

$$\mathcal{TOP}(Q \# Q') = \mathcal{TOP}(Q) \# \mathcal{TOP}(Q') \# (S^2 \times S^{d-1})$$

(B) *If Q is closed but Q' has non-empty boundary then*

$$\mathcal{D}(Q \# Q') = Q \# (-Q) \# \mathcal{D}Q'$$

$$\mathcal{TOP}(Q \# Q') = \mathcal{G}Q \# \mathcal{TOP}(Q')$$

Proof: first observe that for disks D, D' , $(D \# D')$ is diffeomorphic to $D^1 \times S^{d-1}$. Then $Q \# Q'$ is obtained by identifying disks in the boundaries of Q, Q' with corresponding disks in the boundary of $D^1 \times S^{d-1}$. This *is not* the connected sum along the boundary of those three manifolds because not all their boundaries are connected; actually Q and Q' are attached along different components of the boundary of $D^1 \times S^{d-1}$. However, this problem again disappears after multiplying by D^1 : $Q \times D^1$, $Q' \times D^1$ and the intermediate manifold $D^2 \times S^{d-1}$ all have connected boundaries and we have the connected sum along the boundary:

$$(Q \# Q') \times D^1 = (Q \times D^1) \amalg (Q' \times D^1) \amalg (D^2 \times S^{d-1}).$$

Taking the boundary of this manifold and of its product with D^1 we obtain A).

As for B) just observe that $Q \# Q' = Q_{-1} \amalg Q'$ and take products and boundaries.

2.2 Cutting vertices.

Let P be a simple polytope of dimension d and m facets and $P_{\mathbf{v}}$ be a simple polytope of dimension d obtained from P by cutting off one vertex v .

Let $Z_{\mathbf{v}} = Z(P_{\mathbf{v}})$ be the corresponding connected manifold¹⁵

We will show that the operation $Z_{\mathbf{v}}$ preserves connected sums of sphere products. However, the result is not simply connected since $P_{\mathbf{v}}$ is never dual 2-neighborly because the new created facet does not intersect any of the facets not adjacent to v . So it cannot be combined with the theorems of section 1. We can partially avoid this problem by the next construction:

Let $Z_{\mathbf{v}}'$ be the manifold obtained by duplicating the coefficients of the new variable x_0 introduced in the construction of $Z_{\mathbf{v}}$. Notice that this notation is consistent with that of section 1.

Theorem 2.1 :

1.- $Z_{\mathbf{v}}$ is diffeomorphic to $Z \# Z \# (2^{m-d} - 1)(S^1 \times S^{d-1})$.

2.- $Z_{\mathbf{v}}'$ is diffeomorphic to $\mathcal{G}Z \# (2^{m-d} - 1)(S^2 \times S^{d-1})$.

So, if Z is a connected sum of sphere products, any manifold obtained from it by repeatedly applying the operations $Z_{\mathbf{v}}$ and $Z_{\mathbf{v}}'$ is also a connected sum of sphere products.

Proof: We look at Z as the quotient of $P \times \mathbf{Z}_2^m$ by making identifications on the facets of P . Assume that the vertex v lies in facets $F_1, \dots, F_d$ of P and the rest of the facets are $F_{d+1}, \dots, F_m$. Call F_0 the new facet of $P_{\mathbf{v}}$ left by the removal of a small simplex Δ around v .

Then $Z_{\mathbf{v}}$ is obtained from $P_{\mathbf{v}} \times \mathbf{Z}_2^{m+1}$ by making the identifications on all of its facets. Let $Z_{\mathbf{v}+}$ be the manifold with boundary obtained by making all identifications on the facets of $P_{\mathbf{v}} \times \mathbf{Z}_2^m$, except for F_0 . Then, in the notation of section 2.1:

$$Z_{\mathbf{v}} = \mathcal{D}Z_{\mathbf{v}+}$$

$$Z_{\mathbf{v}}' = \mathcal{TOP}(Z_{\mathbf{v}+})$$

Now, $Z_{\mathbf{v}+}$ is Z minus U where U is obtained from $\Delta \times \mathbf{Z}_2^m$ by making the identifications on the subsets $\Delta \cap F_i$, where F_i are the facets of P .

¹⁵This manifold, the manifold $Z_{\mathbf{e}}$ and any other one obtained by cutting a face of Z can be expressed as an intersection of quadrics by intersecting P with a half space that contains all the vertices of P that do not belong to the face, as explained in section 0.2.

If we make the identifications on $\Delta \times \mathbf{Z}_2^d$ on the facets $\Delta \cap F_i$ for $i = 1, \dots, d$ we obtain a disk¹⁶ of dimension d , D^d . On the product $D^d \times \mathbf{Z}_2^{m-d}$ we still have to make identifications on the subsets $\Delta \cap F_i$ for $i = d+1 \dots m$, but these intersections are empty. Therefore $U = D^d \times \mathbf{Z}_2^{m-d}$ and $Z_{\mathbf{V}+}$ is Z with 2^{m-d} disks removed.

Now we use the lemma:

Lemma. *Let M^d be a connected manifold and M_{-k} be obtained from M by removing k open disks. Then*

- 1) $M_{-k} \times D^1 = (M_{-1} \times D^1) \# \Pi(k-1)(D^2 \times S^{d-1})$.
- 2) $\mathcal{D}(M_{-k}) = M \# (-M) \# (k-1)(S^1 \times S^{d-1})$.
- 3) $\mathcal{TOP}(M_{-k}) = \mathcal{G}M \# (k-1)(S^2 \times S^{d-1})$.

Proof: Since M is connected, we can assume all the open disks lie in the interior of a closed disk and therefore M_{-k} can be seen as M_{-1} with $k-1$ copies of $D^1 \times S^{d-1}$ attached along disks in its boundary. This is *not* a connected sum along the boundary, since the boundary of $D^1 \times S^{d-1}$ is not connected, but after multiplying times D^1 this problem disappears and $M_{-k} \times D^1$ is the connected sum claimed in 1). Taking its boundary we get part 2) and taking its boundary after multiplying again by D^1 gives part 3). So the lemma is proved and so are parts 1) and 2) Theorem 2.1.

The last part is obvious for the operation $Z_{\mathbf{V}}$ and it follows from Fico's Lemmata (section 2.1) for the operation $Z_{\mathbf{V}}'$

Remarks:

Although the combinatorial type of $P_{\mathbf{V}}$ depends on which vertex is cut off, the diffeomorphism type of $Z_{\mathbf{V}}$ does not. This gives many examples of different polytopes that produce the same manifold Z .

The analogous result for $Z^{\mathbf{C}}$ is proved in [B-M].

In particular, if P is obtained from a simplex by succesively cutting off vertices then $Z(P)$ is a connected sum of sphere products. Such a polytope

¹⁶ If we had made identification on $\Delta \times \mathbf{Z}_2^{d+1}$ on all the facets we would have obtained the sphere S^d .

is a *dual stack polytope*, and its dual can be obtained from a simplex by successively constructing pyramids on facets.

Its diffeomorphism type depends only on the number of times each operation is applied and not on which vertices, or in which order, they are applied. As an exercise, write the formula for the effect on the manifold Z of successively cutting off n vertices from P .

Part 2 of Theorem 2.1 can be combined with the theorems of section 1 to obtain that connected sums are preserved when applying repeatedly the Z^J and the $Z_{\mathbf{V}}'$ constructions. A more complete version of this will appear as Theorem 2.4 below.

Nevertheless, one would like to have a much more general vertex cutting result. We do not know, for example if $Z^{\mathbf{C}}(P_{\mathbf{V}})$ is a connected sum of sphere products whenever $Z^{\mathbf{C}}(P)$ is.

More generally, we can make the following

Conjecture:

$$Z_{\mathbf{V}}^{\mathbf{C}} \text{ is diffeomorphic to } \mathcal{G}(Z^{\mathbf{C}}) \# \bigg\#_{j=1}^{m-d} \binom{m-d}{j} (S^{j+2} \times S^{d+m-j-1}).$$

This conjecture is consistent with the above theorem and with results of [McG] and [B-M]. We now prove it under a restriction on the number of facets of the polytope:

Theorem 2.2: *If P is polytope of dimension d with $m < 3d$ facets then the above conjecture is true.*

We will follow the same strategy as in the previous results, although now the steps are less elementary:

1) $Z^{\mathbf{C}}(P)$ is obtained from $P \times (S^1)^m$ by making identifications on the facets. To obtain $Z^{\mathbf{C}}(P_{\mathbf{V}})$ we have to multiply $(P_{\mathbf{V}})$ by $(S^1)^{m+1}$ and make the identifications on the $m+1$ facets. Let $Z_+^{\mathbf{C}}(P_{\mathbf{V}})$ be the manifold with boundary obtained from the product $(P_{\mathbf{V}}) \times (S^1)^m$ by making the identifications on all the facets except the new facet 0. Then $Z^{\mathbf{C}}(P_{\mathbf{V}})$ is obtained from $Z_+^{\mathbf{C}}(P_{\mathbf{V}})$ by

multiplying times S^1 and making the identifications on its boundary. In the notation of section 2.1 this means

$$Z^{\mathbf{C}}(P_{\mathbf{V}}) = \mathcal{TOB}(Z_+^{\mathbf{C}}(P_{\mathbf{V}})).$$

So we have to understand $Z_+^{\mathbf{C}}(P_{\mathbf{V}}) \times D^1$. Now $Z_+^{\mathbf{C}}(P_{\mathbf{V}})$ is obtained from $Z^{\mathbf{C}}(P)$ by removing the manifold U obtained from $\Delta^d \times (S^1)^m$ by making the same identifications as for $Z^{\mathbf{C}}(P)$, where Δ^d is the simplex removed to create the new facet 0. If we make the identifications on $\Delta^d \times (S^1)^d$ corresponding to the facets of Δ^d that are part of facets of C we obtain the disk D^{2d} . On the product $\Delta^d \times (S^1)^m$ we still have to make identifications on the intersections with the $m-d$ facets not adjacent to the vertex v but these intersections are empty. Therefore $U = D^d \times (S^1)^{m-d}$ and $Z_+^{\mathbf{C}}(P_{\mathbf{V}})$ is $Z^{\mathbf{C}}(P)$ with a thickened $(m-d)$ -torus removed.

2) This $(m-d)$ -torus can be contracted to a point in $Z^{\mathbf{C}}(P)$ because it is an orbit of the action of T^m (see section 0.2). Since $2(m-d) < m+d$ it is isotopic to an $(m-d)$ -torus inside an open disk in $Z^{\mathbf{C}}(P)$. Therefore

$$Z_+^{\mathbf{C}}(P_{\mathbf{V}}) = Z^{\mathbf{C}}(P) \# (S^{m+d} \setminus (T^{m-d} \times \text{int}(D^{2d}))).$$

Again, since $2(m-d) < m+d$ we can apply Theorem A2.3 of the Appendix to the last summand and Lemma 1 of section 2.1 to the whole sum to obtain the Theorem.


Without the restriction on the number of facets of P , it can be proved that the torus $(m-d)$ -torus can be engulfed by an open disk in $Z^{\mathbf{C}}(P)$. To prove the general conjecture, one would need to prove that this torus is in some sense *standard* inside that disk and some version of Theorem A2.3 of Appendix A3 for that type of *standard* torus.

A more general formula can be conjectured. It includes the case where the basic manifold Z is not a connected sum of sphere products, in the spirit of Theorem 2.1, and where the construction Z^J is immediately applied after cutting off a vertex. Again a partial case can be deduced from Theorem A2.2 of the Appendix.

We need also, of course, a way to include all the non-simply connected manifolds into the picture.

Example: the truncated cube.

We will study now the cube $C = I \times I \times I$ and the truncated cube $C_{\mathbf{V}}$.



Now we have

$$Z(C) = S^1 \times S^1 \times S^1$$

$$Z^{\mathbf{C}}(C) = S^3 \times S^3 \times S^3$$

From Theorem 2.1 we know that

$$Z(C_{\mathbf{V}}) = (S^1 \times S^1 \times S^1) \# (S^1 \times S^1 \times S^1) \# 7(S^1 \times S^2)$$

From Theorem 2.2 we know that

$$Z^{\mathbf{C}}(C_{\mathbf{V}}) = \mathcal{G}(S^3 \times S^3 \times S^3) \# 3(S^3 \times S^7) \# 3(S^4 \times S^5) \# (S^5 \times S^5).$$

It follows from the cohomology ring of $\mathcal{G}(S^3 \times S^3 \times S^3)$ that this manifold does not decompose into a non-trivial connected sum: by the argument of the Lemma in section 3, the 3-dimensional generators would have to lie in one of the summands and so would the rest of the generators of dimension less than 10 since they are related by non-trivial products with the first ones,

so one of the summands must be a homotopy sphere. This shows that we have reached the whole decomposition of $Z^C(C_V)$ as a connected sum.

This answers another question in [B-M], p.111. It will prove useful in section 3.

We can also describe the topology of the truncated d -cube. In fact, Theorem 2.5 gives us the topology of Z^C of an iterated truncation of up to d vertices of the d -cube.

2.3 Cutting edges and other faces.

Let P be a simple polytope of dimension d and m facets and P_e be a simple polytope of dimension d obtained from P by cutting off one edge e .

Let $Z_e = Z(P_e)$. Again, the resulting manifold is not simply connected, but once more we can prove that it becomes simply connected after applying to it the construction Z' based on the new facet of P_e . Let Z_e' be that manifold.

Theorem 2.3: *If Z is 1-connected (or, equivalently, if P is dual 2-neighborly) then*

- 1) Z_e is diffeomorphic to $Z \# Z \# (2^{m-d-1})(S^2 \times S^{d-2}) \# (2^{m-d-1} - 1)(S^1 \times S^{d-1})$.
- 2) Z_e' is diffeomorphic to $\mathcal{G}(Z) \# (2^{m-d-1})(S^3 \times S^{d-2}) \# (2^{m-d-1} - 1)(S^2 \times S^{d-1})$.

Proof: We look at Z as the quotient of $P \times \mathbf{Z}_2^m$ by making identifications on the facets of P . Denote by $F_1, \dots, F_m$ these facets, where $F_1, \dots, F_{d+1}$ are the ones touched by e . Call F_0 the new facet of P_e left by the removal of a regular neighborhood $e \times \Delta^{d-1}$ of e .

Then Z_e is obtained from $P_e \times \mathbf{Z}_2^{m+1}$ by making the identifications on its $m+1$ facets. Let Z_{e+} be obtained from $P_e \times \mathbf{Z}_2^m$, by making identifications on the facets $F_1, \dots, F_m$. Then clearly we have

$$Z_e = \mathcal{D}(Z_{e+})$$

Also, Z_e' is obtained from $P_e \times \mathbf{Z}_2^m \times S^1$ by making the identifications on its facets, so, in the notation of section 2.1,

$$Z_{\mathbf{e}'} = \mathcal{TOP}(Z_{\mathbf{e}+})$$

So we have to describe $Z_{\mathbf{e}+} \times D^1$ and $Z_{\mathbf{e}+} \times D^2$ in order to know their boundaries.

Now $Z_{\mathbf{e}+}$ is Z minus U where U is obtained from $e \times \Delta^{d-1} \times \mathbf{Z}_2^m$ by making the identifications on its intersections with the facets $F_1, \dots, F_m$ of P .

If we make first the identifications on $e \times \Delta^{d-1} \times \mathbf{Z}_2^{d+1}$ on all its facets we obtain the product $S^1 \times S^{d-1}$, but if we only do it on those facets that are part of the facets of P (i.e., excluding the facet 0) we obtain $S^1 \times D^{d-1}$. On the product $e \times \Delta^{d-1} \times \mathbf{Z}_2^m$ we still have to make identifications on the intersections with the facets i for $i = d+2, \dots, m$ but these intersections are empty. Therefore $U = S^1 \times D^{d-1} \times \mathbf{Z}_2^{m-d-1}$ and $Z_{\mathbf{e}+}$ is Z with 2^{m-d-1} copies of $S^1 \times D^{d-1}$ removed.

Then we have the lemma¹⁷:

Lemma. *Let M^d be a simply connected manifold of dimension at least 4 and $M_{\sim k}$ be obtained from M by removing $k > 0$ copies of $S^1 \times D^{d-1}$.*

Then $M_{\sim k} \times D^1$ is diffeomorphic to

$$M_{-1} \times D^1 \amalg k(D^3 \times S^{d-2}) \amalg (k-1)(D^2 \times S^{d-1}).$$

Proof: Since M is simply connected, we can assume all copies of $S^1 \times D^{d-1}$ lie inside an open disk, so we have only to prove it for $M = S^d$.

In this case the result is true for $k = 1$ (even before multiplying by D^1) since S^d minus a copy of $S^1 \times D^{d-1}$ is diffeomorphic to $D^2 \times S^{d-2}$.

Inductively, $M_{\sim(k+1)}$ is obtained from $M_{\sim k}$ by removing from it one copy of $S^1 \times D^{d-1}$ so $M_{\sim(k+1)} = M_{\sim k} \# M_{\sim 1}$.

Now Lemma 2 of section 2.1 gives the induction step in the case $M = S^d$:

$$\begin{aligned} M_{\sim(k+1)} \times D^1 &= (M_{\sim k} \# (M_{\sim 1} \times D^1)) = (M_{\sim k} \times D^1) \amalg (M_{\sim 1} \times D^1) \amalg (D^2 \times S^{d-1}) = \\ &= (M_{-1} \times D^1) \amalg k(D^3 \times S^{d-2}) \amalg (k-1)(D^2 \times S^d) \amalg (D^3 \times S^{d-2}) \amalg (D^2 \times S^{d-1}) \\ &= (M_{-1} \times D^1) \amalg (k+1)(D^3 \times S^{d-2}) \amalg k(D^2 \times S^d). \end{aligned}$$

and the Lemma is proved.

¹⁷Recall the notation of section 2.1: M_{-1} denotes M minus an open disk. For dimension 3 the lemma is not true, but the theorem is, since the only possible polytope is the simplex.

Applying it to Z we obtain $Z_{\mathbf{e}+} \times D^1$ is diffeomorphic to

$$(Z_{-1} \times D^1) \amalg 2^{m-d-1}(D^3 \times S^{d-2}) \# (2^{m-d-1} - 1)(D^2 \times S^{d-1}).$$

Taking its boundary we obtain 1). Taking the boundary of its product with D^1 we obtain 2).

Remarks:

1.- Although the combinatorial type of $P_{\mathbf{e}}$ depends on which edge is cut off, the diffeomorphism type of $Z_{\mathbf{e}}$ does not. This gives many new examples of different polytopes that produce the same manifold Z .

2.- If Z is not simply connected then Theorem 2.6 and the previous observation are not true. For example, if P is the triangular prism $\Delta^2 \times I$, then by cutting off an edge we can obtain either the pentagonal book or the cube, whose corresponding manifolds are $\#5(S^1 \times S^2)$ and $S^1 \times S^1 \times S^1$, respectively. The reason lies in the fact that when the edge connects two facets of P that do not intersect, then this edge produces an embedded S^1 that is not trivial in $H_1(Z)$ and therefore cannot be contained inside a disk. It remains to be clarified what happens in this case.

3.- One problem with part 1 is that it cannot be iterated since the resulting manifold $Z_{\mathbf{e}}$ is not simply connected.

Combining Theorem 2.3 with our previous theorems we get

Theorem 2.4: *If $Z = Z(P)$ is a connected sum of sphere products and is simply connected of dimension at least 5, then any manifold obtained from Z by repeatedly applying the Z^J , $Z_{\mathbf{v}}'$ and $Z_{\mathbf{e}}'$ constructions (any number of times each and in any order) as well as its corresponding moment-angle manifold, is also a connected sum of sphere products.*

The basic manifold can be any sphere, any $Z(P)$ with free homology for a dual-neighborly polytope or an adequate Z^J for a polygon (Theorems 1.3 and 1.4) or any simply connected manifold with $m - d = 3$ of dimension at least 5 that is not a triple sphere product ([LdM1]).

So this theorem seems to include all cases known so far of moment-angle manifolds that are connected sums of sphere products.

Remarks: The theorems on this section can be generalized to the process of removing any simplicial face of P but need, of course, stronger connectivity hypotheses.

One can conceive also results about removing non-simplicial faces of P . It has been conjectured that any simple polytope can be realized as a face of a dual neighborly polytope, so, in principle we need to consider the effect of cutting off any possible simple polytope.

It is known that there are numerous dual neighborly polytopes in each dimension. If to these we add the infinite family stemming from each of them by applying the three constructions above (which produce polytopes which are *not* dual neighborly) we can conclude that there is a large number of polytopes P for which $Z(P)$ is a connected sum of sphere products.

3 On the cohomology ring of $Z(P)$.

In [B-M] the cohomology ring of $Z^C(P)$ is described (by an explicit rule) in terms of intersection products on the relative homology groups of pairs (P, P_J) where P_J is a union of facets of P . A slightly different rule, valid for general moment-angle complexes, appears in [Ba]. We shall see now by examples that these rules must be drastically modified in the case of the manifolds $Z(P)$.

We will show that the cohomology rings of the manifolds associated to the truncated cube are not isomorphic as *ungraded rings*.

We will consider the cohomology rings with $\mathbf{Z}_2$ coefficients. Since all the homology groups of the spaces involved are free, this will be enough to establish the difference without having to compute signs, so in fact we have an stronger statement.

We have shown at the end of section 2.2 that

$$Z(C_V) = (S^1 \times S^1 \times S^1) \# (S^1 \times S^1 \times S^1) \# 7(S^1 \times S^2)$$

$$Z^C(C_V) = \mathcal{G}(S^3 \times S^3 \times S^3) \# 3(S^3 \times S^7) \# 3(S^4 \times S^5) \# (S^5 \times S^5).$$

We begin by describing the cohomology ring of the first summands of each manifold:

$$X := (S^1 \times S^1 \times S^1) \# (S^1 \times S^1 \times S^1)$$

$$Y := \mathcal{G}(S^3 \times S^3 \times S^3)$$

$H^*(X)$ has:

One generator 1 in dimension 0.

6 generators $a_1, a_2, a_3, a'_1, a'_2, a'_3$ in dimension 1.

6 generators $b_{1,2}, b_{1,3}, b_{2,3}, b'_{1,2}, b'_{1,3}, b'_{2,3}$ in dimension 2.

One generator f (the fundamental class) in dimension 3.

The multiplication rules are:

$$a_i a_j = b_{i,j} \text{ for } i \neq j$$

$$a_1 a_2 a_3 = t$$

$$a'_i a'_j = b'_{i,j} \text{ for } i \neq j$$

$$a'_1 a'_2 a'_3 = f$$

all other products not derived from these being 0.

On the other hand $H^*(Y)$ has:

One generator 1 in dimension 0.

3 generators A_1, A_2, A_3 in dimension 3.

3 generators A'_1, A'_2, A'_3 in dimension 4.

3 generators $B_{1,2}, B_{1,3}, B_{2,3}$ in dimension 6.

3 generators $B'_{1,2}, B'_{1,3}, B'_{2,3}$ in dimension 7.

One generator F' (the fundamental class) in dimension 10.

(In this case a generator is denoted by C' when it is inherited from the element $C \times u$ in the cohomology of $(S^3 \times S^3 \times S^3) \times S^1$ where u is the generator of $H^1(S^1)$).

The multiplication rules are:

$$A_i A_j = B_{i,j} \text{ for } i \neq j$$

$$A_i A'_j = B'_{i,j} \text{ for } i \neq j$$

$$A_i A_j A'_k = F' \text{ for } i, j, k \text{ all different.}$$

all other products not derived from these being 0.

Observe in particular that

$$A'_i A'_j = 0 \quad (*)$$

for all i, j , simply because the product lies in $H^8(Y) = 0$.

This is the main difference with the previous ring: three indecomposable, linearly independent elements have all their products equal to zero. To make this idea into a formal proof of non-isomorphism we have to work modulo the decomposable elements:

In both rings there is the maximal ideal $\mathcal{M}$ of non-units, spanned by all the generators except 1. The square of this ideal $\mathcal{M}^2$ is generated by the decomposable elements and its cube is generated by the fundamental class. There is an induced multiplication

$$M : \mathcal{M}/\mathcal{M}^2 \times \mathcal{M}/\mathcal{M}^2 \rightarrow \mathcal{M}^2/\mathcal{M}^3$$

An ungraded ring isomorphism between the two rings would have to preserve the above ideals and multiplication.

Proposition. *For the ring $H^*(X)$, any three elements in $\mathcal{M}/\mathcal{M}^2$ all whose products are zero in $\mathcal{M}^2/\mathcal{M}^3$ must be linearly dependent.*

Proof: Let

$$x_j = \sum_{i=1}^3 \lambda_j^i a_i + \sum_{i=1}^3 \mu_j^i a'_i$$

for $i = 1, \dots, 3$ be (representatives of) such elements. Then

$$x_j x_k = \sum_{r < s} (\lambda_j^r \lambda_k^s + \lambda_j^s \lambda_k^r) b_{r,s} + \sum_{r < s} (\mu_j^r \mu_k^s + \mu_j^s \mu_k^r) b'_{r,s}.$$

The fact that this product is 0 means that all 2×2 determinants of the matrices λ_j^i and μ_j^i are 0. This means that in each there is at most one linearly independent column and that the matrix of coefficients of the x_i , which is formed by the columns of those two, has rank at most 2. Thus proposition 4.1 is proved.

To complete the rings of $Z(C_V)$ and $Z^C(C_V)$ one has to add in each case to the corresponding previous one 7 couples of elements u_i, v_i such that each product $u_i v_i$ is the fundamental class and the product of each u_i, v_i with all the rest of generators is 0. : We have now¹⁸

¹⁸An analogous result would be true for the truncated d -dimensional cube for $d > 2$, but not for $d = 2$ since the manifolds of the pentagon have the same ungraded cohomology ring. Observe how the above proof fails in this case.

Theorem 3.1 *The rings $H^*(Z(C_V), \mathbf{Z}_2)$ and $H^*(Z^C(C_V), \mathbf{Z}_2)$ are not isomorphic as ungraded rings..*

Proof: We follow the same lines as for the previous proposition: now the rings considered there must be complemented with 7 couples of generators u_i, v_i such that each product $u_i v_i$ is the fundamental class and the product of each u_i and v_i with all the rest of generators is 0. Since the fundamental class is in $\mathcal{M}^3$ this means that $H^*(Z^C(C_V))$ has 17 generators in $\mathcal{M}/\mathcal{M}^2$ all whose products are 0 under the induced multiplication M . If in the case of $H^*(Z(C_V), \mathbf{Z}_2)$ we have 17 elements in $\mathcal{M}/\mathcal{M}^2$ for which all products under M are 0, we can express them in terms of the 20 generators a_i, a'_i, u_i, v_i by a 17×20 matrix. By the same computation as above, at most two of the first six columns of this matrix are independent, so the matrix has at most rank 16 and the 17 elements are not linearly independent. This proves the Theorem.

Therefore the product rules in [Ba], [B-M] are not valid in general for the rings $H^*(Z(P))$, even if the necessary adjustments to the degrees were done.

It can be shown that the cohomology rings of $Z^C(P)$ and $Z^C(P')$ are isomorphic as *ungraded rings*. (See [B-B-C-G-LdM]); this should be also a consequence of the other known descriptions of the cohomology rings, [B-M], [B-P], [D]). For example, if P is a product of two simplices then $Z^C(P) = S^{2p-1} \times S^{2q-1}$ and $Z^C(P') = S^{2p+1} \times S^{2q-1}$ and the ungraded cohomology rings are isomorphic, with generators $\{1, x, y, xy\}$. Or, if $Z^C(P)$ is a connected sum of sphere products then $Z^C(P')$ is also one, with the same number of summands, but where inside each summand one of the factors has risen its dimension by two.

The previous Theorem implies that this also cannot be true in general for the cohomology rings of $Z(P)$ and $Z(P')$, not even when taken with $\mathbf{Z}_2$ coefficients, since the passage from $Z(P)$ to $Z^C(P)$ goes through the iteration of the operation $Z(P')$.

If L is a collection of facets of P , we denote by P_L the union of the facets of P in L .

We claim that the correct rules are as follows:

Intersection product: There is an isomorphism

$$H_i(Z) \approx \bigoplus_L H_i(P, P_L)$$

so that the intersection of two homology classes in Z is given by the sum of the intersection products on the right hand side

$$H_i(P, P_L) \otimes H_j(P, P_J) \rightarrow H_{i+j-d}(P, P_{L \cap J}).$$

Cup product: There is an isomorphism

$$H^i(Z) \approx \bigoplus_L H^i(P, P_L)$$

so that the cup product of two cohomology classes in Z is given by the sum of the cup products on the right hand side

$$H^i(P, P_L) \otimes H^j(P, P_J) \rightarrow H^{i+j}(P, P_{L \cup J}).$$

In other words, the rule that a product must be 0 unless L and J satisfy a strict set-theoretical condition, must be dropped.

Details and extensions to generalized moment-angle complexes will appear in a forthcoming article ([B-B-C-G-LdM]).

Appendix.

A1. Recognizing a connected sum of sphere products.

Recall that, after Thom, Smale and Milnor, the best way to identify a compact manifold M is to study an adequate compact manifold Q whose boundary is M . For example, if M^n is a homotopy sphere, we cannot tell if it is diffeomorphic to the standard sphere S^n by looking at the usual topological invariants of M : since these are homotopy invariants there is nothing to look at. But we know that M is the boundary of some manifold Q . The first thing to ask is if Q is paralellizable. If there is no such paralellizable Q then M is not standard since S^n bounds a disk. If one finds a Q that is paralellizable,

then one can decide if M is standard by studying topological invariants of Q ([K-M]).

Now a connected sum of sphere products

$$\#_{i=1}^k (S^{p_i} \times S^{d-p_i})$$

is the boundary of the connected sum along the boundary

$$\coprod_{i=1}^k (S^{p_i} \times D^{d-p_i+1})$$

which has the following properties: It is simply connected if $1 < p_i$, has simply connected boundary if $1 < p_i < d - 2$, its homology groups are free and bases of them can be represented by disjoint embedded spheres with trivial normal bundle.

To determine such a connected sum it is enough to verify those conditions:

Theorem A1. *Let Q be a manifold with boundary satisfying:*

- a) Q is of dimension $d + 1 \geq 6$
- b) Q is simply connected with simply connected boundary.
- c) $H_i(Q)$ is free with basis $\alpha_{i,j}$ for all i and $H_i(Q) = 0$ for $i \geq d - 1$
- d) *There is a collection $\{S_j^i\}$ of disjoint embedded spheres with trivial normal bundle inside Q that represent basis elements $\alpha_{i,j}$ of $H_i(Q)$.*

Then Q is diffeomorphic to a connected sum along the boundary

$$\coprod_{i=1}^k (S^{p_i} \times D^{d-p_i+1})$$

and therefore ∂Q is diffeomorphic to

$$\#_{i=1}^k (S^{p_i} \times S^{d-p_i}).$$

Proof: For every sphere S_j^i take a closed product neighborhood $U_{i,j} = S_j^i \times D^{d-i}$ in such a way that they are all disjoint. Connect all $U_{i,j}$ by a minimal collection of thin tubes and call Q_1 the union of all the $U_{i,j}$ and

their connecting tubes, which we can assume is contained in the interior of Q . Then Q_1 is a connected sum of the required form for Q and we will prove that these two manifolds are diffeomorphic. Let H be equal to Q minus the interior of Q_1 , so that H is a cobordism between the manifolds ∂Q and ∂Q_1 .

Now Q_1 , ∂Q_1 are simply connected because all the spheres S_j^i have dimension $2 \geq i \geq d - 2$ and so is H because Q is simply connected and every loop in H bounds a disk in Q that can be made disjoint from Q_1 since the spheres have codimension greater than 2. Now $H_i(H, \partial Q_1) = H_i(Q, Q_1) = 0$ for all i because the inclusion of Q_1 in Q induces isomorphisms of homology groups. Therefore H is a simply connected h -cobordism of dimension at least 6 between simply connected manifolds and so is a product by the h -cobordism theorem. This proves that Q is diffeomorphic to Q_1 and the theorem is proved.

A2. Complements of sphere products in spheres.

We will work in the unit sphere S^n in $\mathbf{R}^{n+1}$ and assume we have a decomposition

$$\mathbf{R}^{n+1} = \mathbf{R}^{n_1+1} \times \dots \times \mathbf{R}^{n_k+1} \times \mathbf{R}^q.$$

We have then $n = \sum_{i=1}^k n_i + q + k - 1$ and we will denote an element in $\mathbf{R}^{n+1}$ as a $(k+1)$ -tuple $(X_1, \dots, X_k, Y)$ with $X_i \in \mathbf{R}^{n_i+1}$ and $Y \in \mathbf{R}^q$.

We will denote by $\{e_j^i\}$ for $j = 0, \dots, n_i$ the standard basis of $\mathbf{R}^{n_i+1}$.

We will consider the product of spheres

$$\mathcal{P} = S^{n_1} \times \dots \times S^{n_k} \times \{0\}$$

given by the equations

$$X_i \cdot X_i = \frac{1}{k}, \quad Y = 0.$$

$\mathcal{P}$ has dimension $\sum_{i=1}^k n_i$ and codimension $q + k - 1$.

Within each sphere S^{n_i} we consider the base point $p_i = \frac{1}{\sqrt{k}} e_0^i = (\frac{1}{\sqrt{k}}, 0, \dots, 0)$.

We are interested in the complement of an open tubular neighborhood of $\mathcal{P}$ in S^n . We will denote this complement by $\mathcal{E}$.

Let $I = \{1, \dots, k\}$. If $J \subset I$ we denote by $|J|$ the number of elements of J and by n_J the sum $\sum_{i \in J} n_i$.


We will make use of the subproduct $\mathcal{P}_J$ of $\mathcal{P}$ defined by the equations

$$\begin{aligned} X_i &= p_i \quad i \notin J \\ Y &= 0. \end{aligned}$$

So $\mathcal{P}_J$ is diffeomorphic to the product $\prod_{i \in J} S^{n_i}$ of dimension n_J , the rest of the coordinates having fixed values. If we change a base point p_i by any other point $p'_i \in S^{n_i}$ we obtain a diffeomorphic product $\mathcal{P}'_J$ that is isotopic to $\mathcal{P}_J$ inside $\mathcal{P}$ if $n_i > 0$.

When $n_i > 0$ for $i = 1, \dots, k$ $\mathcal{P}$ is connected and its reduced homology groups are free with a generator in dimension $n_J > 0$ represented by the fundamental class of $\mathcal{P}_J$ for every non-empty $J \subset I$.

We will show that if $n_i > 0$ for $i = 1, \dots, k$ we have Alexander dual generators for the homology of $\mathcal{E}$ represented by embedded round spheres. The Alexander dual of a single sphere S^{n_i} will be a great sphere of dimension $n - n_i - 1$ lying in a complementary subspace. The Alexander dual of the products of spheres will be spheres that become smaller and more slanted as the number of factors grows. The Alexander dual of the whole product $\mathcal{P}$ will be a small sphere which is a fiber of its spherical normal bundle. A diagrammatic representation of the construction is the following:



Here we have the picture for $n = 2$, $k = 3$, $n_1 = n_2 = 0$ (a case *not* covered by the hypothesis of Theorem A2.1 below). The product $\mathcal{P} = S^0 \times S^0 \times S^0$ is given by the 8 marked points. The vertical great circle is the Alexander dual of an S^0 given by one point on each vertical hemisphere. The smaller vertical circle is the Alexander dual of an $S^0 \times S^0$ consisting of the points in the upper hemisphere since the small disk this circle bounds contains exactly one of its points. The smallest slanted circle is the Alexander dual of the whole product, since it surrounds just one of its points.

Theorem A2.1. *Assume $n_i > 0$ for $i = 1, \dots, k$. Then there is a basis for the homology of $\mathcal{E}$ represented by embedded round spheres. In each dimension it is the Alexander dual basis of the basis $\mathcal{P}_J$ of the reduced homology of $\mathcal{P}$.*

Proof: The result is well-known for $k = 1$ so we assume $k > 1$. We claim that the Alexander dual basis of $\mathcal{P}_J$ in $\mathcal{P}$ is represented by the subspheres S_J of S^n given by the equations

$$X_i = x_i e_0^i, \quad i \in J$$

$$\sum_{i \in J} x_i = \frac{|J| - 1}{\sqrt{k}}.$$

Clearly S_J is a round sphere, being the transversal intersection of S^n with an affine space of codimension $n_J + 1$.

To see that S_J lies in $\mathcal{E}$ just observe that a point in the intersection $S_J \cap \mathcal{P}$ must satisfy for $i \in J$ both $X_i = x_i e_0^i$ and $X_i \cdot X_i = \frac{1}{k}$, so $x_i = \frac{\pm 1}{\sqrt{k}}$. But this implies that

$$\sum_{i \in J} x_i = \frac{q}{\sqrt{k}}.$$

where q is congruent to $|J|$ modulo 2, which is incompatible with the last equation for S_J and the intersection is empty.

Now, S_J bounds inside S^n the disk D_J of codimension n_J given by:

$$X_i = x_i e_0^i, \quad i \in J$$

$$\sum_{i \in J} x_i \geq \frac{|J| - 1}{\sqrt{k}}.$$

A point in $D_J \cap \mathcal{P}_J$ satisfies as above $x_i = \frac{\pm 1}{\sqrt{k}}$ and, if some sign is $-$, then

$$\sum_{i \in J} x_i \leq \frac{|J| - 2}{\sqrt{k}}$$

which is incompatible with the above inequality. From the equations of S_J we obtain that

$$X_i = p_i$$

also for $i \notin J$ and that $Y = 0$.

Therefore the only point in $D_J \cap \mathcal{P}_J$ is

$$p = \sum_{i=1}^k \frac{1}{\sqrt{k}} e_0^i = (p_1, \dots, p_k, 0).$$

We have to verify that the intersection is transversal: the unit normal vector to S^n at p is $\sum_{i=1}^k e_0^i$, so a tangent vector v to S^n at p is characterized by having the sum of its e_0^i components equal to zero. We can decompose v into a sum $v_1 + v_2$ where v_1 is the projection of v on the subspace of $\mathbf{R}^n$ generated by the vectors e_j^i where $i \in J$ and $j > 0$. Such a vector e_j^i is tangent to the sphere S^{n_i} at the point p_i , so v_1 is tangent to the product $\mathcal{P}_J$ at p . Now v_2 has all those components equal to zero and the sum of its e_0^i components

equal to zero. But this fact clearly characterizes the tangent vectors to D_J at p . So v_2 is tangent to D_J at p and therefore the intersection of $\mathcal{P}_J$ and D_J is transversal at p since the (direct) sum of the two tangent spaces is the whole tangent space of S^n . So the linking number of $\mathcal{P}_J$ and S_J is ± 1 .

To end the proof of the Theorem we have to show that $\{\mathcal{P}_J\}$ is the Alexander dual basis of $\{S_J\}$ dimension by dimension. That is, that the linking number of $\mathcal{P}_J$ with S_L is zero for any $L \neq J$ with $n_J = n_L$:

First observe that if $J \subset L$ they would be equal since $n_J = n_L$ and by hypothesis every n_i is positive. Then there is an $i \in J$ not in L . Therefore we have for a point in the intersection $D_J \cap \mathcal{P}_L$ both $X_i = p_i$ and $X_i = x_i e_0^i$. These conditions are not incompatible (in fact, the intersection can have positive dimension in some cases). But if we perturb slightly $\mathcal{P}_L$ so that the first equation becomes $X_i = a e_0^i + b e_1^i$ for $b > 0$ close to 0 and $a^2 + b^2 = 1/k$ (this is possible since n_i is positive) we obtain a new product $\mathcal{P}'_L$ isotopic to $\mathcal{P}_L$ such that $D_J \cap \mathcal{P}'_L = \emptyset$. Since this new product represents the same homology class as $\mathcal{P}_L$ we see that the intersection number is zero and the Theorem is proved.

For $k > 1$ we cannot apply to $\mathcal{E}$ Theorem A.1 characterizing a connected sum along the boundary (and in fact it is not, since its boundary is a product of at least three spheres). But we can apply it to $\mathcal{E} \times D^1$ since the spheres S_J can be embedded with different values of the coordinate in D^1 so they are disjoint, represent a basis of its homology and have codimension at least 3 and dimension at least 2 if $q + k \geq 4$. As a consequence we have:

Theorem A2.2. *If $q + k \geq 4$, then*

A) $\mathcal{E} \times D^1$ is diffeomorphic to

$$\coprod_{\emptyset \neq J \subset I} (S^{n-n_J-1} \times D^{n_J+2}).$$

B) The double of $\mathcal{E}$, $\mathcal{D}(\mathcal{E})$, is diffeomorphic to

$$\sharp_{\emptyset \neq J \subset I} (S^{n-n_J-1} \times S^{n_J+1}).$$

C) The trivial open book on $\mathcal{E}$, $\mathcal{TOP}(\mathcal{E})$, is diffeomorphic to

$$\sharp_{\emptyset \neq J \subset I} (S^{n-n_J-1} \times S^{n_J+2}).$$

When $n_i > 0$ for $i = 1, \dots, k$, A) follows from Theorem A1 since the hypothesis $q + k \geq 4$ implies that $\mathcal{P}$ has codimension at least 3 in S^n and so $\mathcal{E} \times D^1$ is simply connected with simply connected boundary. When some n_i are zero it follows by induction on the number of them: if we add a new factor to a product $\mathcal{P}$ we get a new complement $\mathcal{E}'$ which is clearly the connected sum of two copies of $\mathcal{E}$. By Lemma 1.2, $\mathcal{E}' \times D^1$ is the connected sum along the boundary of two copies of $\mathcal{E} \times D^1$ and one copy of $S^{n-1} \times D^2$ and by the induction hypothesis the collection of summands of the form $S^{n-n_j-1} \times D^{n_j+2}$ coincides with the one claimed for $\mathcal{E}' \times D^1$ and A) is proved. B) follows from A) by taking its boundary and C) by taking its product with D^1 and again the boundary.

In the special case where all $n_i = 1$ (i.e., when $\mathcal{P}$ is the k -torus) this result takes the form

Theorem A2.3. *Consider an embedding of the torus T^k in S^n with $2k < n$ and $\mathcal{E}$ the complement of an open tubular neighborhood of T^k . Then*
A) $\mathcal{E} \times D^1$ is diffeomorphic to

$$\prod_{j=1}^k \binom{k}{j} (S^{n-j-1} \times D^{j+2}).$$

B) The double of $\mathcal{E}$, $\mathcal{D}(\mathcal{E})$, is diffeomorphic to

$$\prod_{j=1}^k \binom{k}{j} (S^{n-j-1} \times S^{j+1}).$$

C) The trivial open book on $\mathcal{E}$, $\mathcal{TOP}(\mathcal{E})$, is diffeomorphic to

$$\prod_{j=1}^k \binom{k}{j} (S^{n-j-1} \times S^{j+2}).$$

This follows from the previous theorem since any embedding of the torus T^k in S^n with $2k < n$ is isotopic to the product considered there.

Part B) was proved by McGavran in the PL-category ([Mc]) but for a more general situation: A *standard torus* lying in a sphere with any codimension ≥ 3 . It does not seem that analogs of the other parts of the theorem can be deduced easily from his results.

References

- [A-LL] D. Allen, and J. La Luz, *A Counterexample to a conjecture of Bosio and Meersseman*, Contemporary Mathematics 460, AMS, 2008, 37-46.
- [B-B-C-G], A. Bahri, M. Bendersky, F. R. Cohen, and S. Gitler, *The polyhedral product functor: a method of computation for moment-angle complexes, arrangements and related spaces*, arXiv:0711.4689v2 [math.AT] 8 Dec 2008.
- [B-B-C-G2], A. Bahri, M. Bendersky, F. R. Cohen, and S. Gitler, *An infinite family of toric manifolds associated to a given one*, manuscript.
- [B-B-C-G-LdM], A. Bahri, M. Bendersky, F. R. Cohen, S. Gitler and S. López de Medrano, *Cup products in generalized moment angle complexes*, in preparation.
- [Ba1] I.V. Baskakov, *Cohomology of K -powers of spaces and the combinatorics of simplicial subdivisions* Russian Math. Surveys 57(2002) 989-990.
- [Ba2] I.V. Baskakov, *Massey triple products in the cohomology of moment-angle complexes*. Russian Math. Surveys 58 (2003), no.5, 1039-1041.
- [B], F. Bosio, *Variétés complexes compactes: une généralisation de la construction de Meersseman et López de Medrano-Verjovsky*. Ann. Inst. Fourier (Grenoble), 51 (2001), 1259-1297.
- [B-M], F. Bosio and L. Meersseman, *Real quadrics in $\mathbf{C}^n$, complex manifolds and convex polytopes*. Acta Math. 197 (2006), no. 1, 53-127.
- [B], W. Browder, *Surgery on simply-connected manifolds*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 65. Springer-Verlag, New York-Heidelberg, 1972. ix+132 pp.
- [B-P], V.M. Buchstaber and T.E. Panov, *Torus actions and their applications in Topology and Combinatorics*, University Lecture Series, AMS (2002).
- [Ch], M. Chaperon, *Géométrie différentielle et singularités de systèmes dynamiques*. Astérisque 138139 (1986), 434 pages.
- [Ch-LdM], M. Chaperon and S. López de Medrano, *Birth of attracting compact invariant submanifolds diffeomorphic to moment-angle manifolds in*

generic families of dynamics, C. R. Acad. Sci. Paris, Ser. I 346 (2008) 1099-1102.

[dL], M. de Longueville, *The ring structure on the cohomology of coordinate subspace arrangements*, Math. Z. 233 (2000) 553-577.

[D-S], G. Denham, A. Suciu, *Moment-angle complexes, monomial ideals, and Massey products*, Pure and Applied Math. Quarterly 3 (2007), no. 1, 25-60.

[D-J], M. Davis and T. Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. Journal 62 (1991), 417-451.

[GA], F. González Acuña, *Open Books*, University of Iowa Notes, 12 p.

[H], F. Hirzebruch, private conversation, based on *Arrangements of lines and algebraic surfaces*, in Arithmetic and Geometry, Vol. II (= Progr. Math. 36), Boston: Birkhauser, 1983, 113140.

[K-M] M. Kervaire and J. Milnor, *Groups of Homotopy Spheres: I* Annals of Mathematics, 2nd Ser., Vol. 77, No. 3. (1963), pp. 504-537.

[L-M] J. J. Loeb and M. Nicolau, *On the complex geometry of a class of non-Kaehlerian manifolds*. Israel J. Math. 110 (1999), 371-379.

[LdM1], S. López de Medrano, *Topology of the intersection of quadrics in $\mathbf{R}^n$* , in *Algebraic Topology* (Arcata Ca, 1986), Springer Verlag LNM **1370**(1989), pp. 280-292, Springer Verlag.

[LdM2], S. López de Medrano, *The space of Siegel leaves of a holomorphic vector field*, in Holomorphic Dynamics (Mexico, 1986), Lecture Notes in Math., 1345, pp. 233-245. Springer, Berlin, 1988.

[LdM3], S. López de Medrano, *Intersection of quadrics and the polyhedral product functor*, in preparation.

[LdM-V], S. López de Medrano and A. Verjovsky, *A new family of complex, compact, nonsymplectic manifolds*. Bol. Soc. Brasil. Mat., 28 (1997), 253-269.

[Mc] D. McGavran, *Adjacent connected sums and torus actions*, Trans. AMS 251 (1979) 235-254

[Me], L. Meersseman, *A new geometric construction of compact complex manifolds in any dimension*. Math. Ann., 317 (2000), 79115.

[Me-V1], L. Meersseman and A. Verjovsky, *Holomorphic principal bundles over projective toric varieties*. J. Reine Angew. Math., 572 (2004), 5796.

[Me-V2], L. Meersseman and A. Verjovsky, *Sur les variétés LV-M*, Contemporary Mathematics 475, AMS, 2008, 111-134.

[W1], C.T.C. Wall, *Classification of $(n - 1)$ -connected $2n$ -manifolds*. Ann. of Math. (2) 75 1962 163–189.

[W2], C.T.C. Wall, *Classification problems in differential topology. VI. Classification of $(s - 1)$ -connected $(2s + 1)$ -manifolds*. Topology 6 1967 273–296.

[W3], C.T.C. Wall, *Stability, pencils and polytopes*, Bull. London Math. Soc., 12 (1980), 401-421.

sgitler@math.cinvestav.mx

Department of Mathematics, Cinvestav, Mexico.

santiago@matem.unam.mx

Instituto de Matemáticas, Universidad Nacional Autónoma de México.